

R-Parity violating flavor symmetries, recent neutrino data and absolute neutrino mass scale

Gautam Bhattacharyya¹, Heinrich Päs², and Daniel Pidt²

¹⁾ *Saha Institute of Nuclear Physics, 1/AF Bidhan Nagar, Kolkata 700064, India*

²⁾ *Fakultät für Physik, Technische Universität Dortmund, D-44221, Dortmund, Germany*

Abstract

We study the rôle of a very general type of flavor symmetry in controlling the strength of *R*-parity violation in supersymmetric models. We assume that only leptons are charged under a global symmetry whose breaking induces lepton number (and, hence, *R*-parity) violation. The charge assignments of leptons under this symmetry are such that the total number of independent lepton number violating couplings is reduced from 39 to 6. The most severe constraints on these flavor-correlated couplings arise from neutrino masses and mixing as well as from the non-observation of $K_L \rightarrow \mu e$. We find that such a scenario predicts an almost vanishing smallest neutrino mass eigenvalue, allowing the upcoming generation of neutrinoless double beta decay experiments to shed light on the hierarchy.

I Introduction

R-parity in supersymmetry is a discrete symmetry which is defined as $R_p = (-1)^{3B+L+2S}$, where B , L and S are the baryon number, lepton number and spin of the particle respectively (see [1, 2, 3, 4]). All standard model (SM) particles have $R_p = 1$, while all superparticles have $R_p = -1$. The assumption of *R*-parity conservation in supersymmetric models is quite *ad hoc*, as this is not supported by any deep underlying principle. Historically, it was imposed to keep the proton stable. However, proton decay requires a simultaneous presence of B and L violation. Therefore, dropping all R_p violating (RpV) couplings in one go is certainly an overkill [5]. Still, in conventional supersymmetric theories R_p conservation is imposed primarily for the sake of convenience, as otherwise the number of independent parameters in the minimal supersymmetric standard model (MSSM), which is already very large and difficult to handle, is augmented by a set of new RpV parameters. Moreover, conserved R_p implies that the lightest supersymmetric particle is stable, which leads to a plausible dark matter candidate, and also attributes supersymmetry with a characteristic missing energy signature in colliders. On the other hand, if lepton number is violated, one distinct advantage is that neutrino masses can be generated via a perfectly renormalizable interaction [6, 7, 8] without the need of introducing any right-handed neutrinos.

The most general superpotential with explicit RpV couplings is given by [3]

$$W_{RPV} = \mu_i L_i H_u + \frac{1}{2} \lambda_{ijk} L_i L_j E_k^C + \lambda'_{ijk} L_i Q_j D_k^C + \frac{1}{2} \lambda''_{ijk} U_i^C D_j^C D_k^C, \quad (1)$$

where i , j and k are the three quark and lepton generation indices. Here, Q_i and L_i are SU(2)-doublet quark and lepton superfields, respectively; D_i^C and E_i^C are SU(2)-singlet superfields for down-type quarks and charged leptons, respectively; and, H_u is the Higgs superfield that generates the mass of the up-type quarks. This introduces 48 new couplings: 3 μ_i -type, 9 λ_{ijk} -type (note the antisymmetry in the first two indices), 27 λ'_{ijk} -type and 9 λ''_{ijk} -type couplings (note the antisymmetry in the last two indices). Only the λ'' couplings are B violating, the rest are all L violating. Besides, new RpV soft terms appear which introduce more unknown parameters. We do not explicitly write down these soft terms but will mention about the relevant ones in

appropriate places. Dealing with so many new parameters substantially reduces the predictivity of the model. At this point there are two ways to proceed. One may either take one or two RpV couplings as non-vanishing at a time and study their implications, or, apply some suitable flavor symmetry to relate one coupling to another [9, 10]. We shall take the latter approach in this paper (see also [11] which does not differentiate between lepton doublet and down-type Higgs superfields for a different approach following a similar spirit). We show that with a simple flavor hypothesis we can bring down the number of totally independent RpV (more specifically, L violating) couplings to only six. These couplings induce neutrino Majorana masses, and if the neutrino mixing matrix is tri-bimaximal (TBM) then the number of independent L violating couplings can be further reduced to four, a scenario which prefers an inverse neutrino mass hierarchy.

II A generic flavor model

We assume that the Yukawa structure leading to the masses and mixing of quarks and charged leptons is fixed by some unspecified global symmetry. This symmetry also ensures baryon number conservation. There is a second global symmetry (X), an abelian horizontal symmetry, which is at the centre of our attention. *Only* leptons are charged under X , such that for each generation i ,

$$Q_X(L_i) = -Q_X(E_i^C). \quad (2)$$

We assume that the Q_X charges of different generations are all positive. The horizontal symmetry is explicitly broken by a small parameter $\varepsilon < 1$, whose charge under X is $Q_X(\varepsilon) = -1$. If the total charge of a given superpotential term is n , then the term is suppressed by ε^n . As an example, if $Q = Z_N$, then the suppression would be $\varepsilon^{n(\text{mod } N)}$ [9].

Now we look at the consequences of Eq. (2) for the 48 RpV couplings of Eq. (1). Since B -number is conserved, all the λ'' couplings vanish right away. Since only leptons are charged under X , it follows that $Q_X(L_i Q_j D_k^C) = Q_X(L_i H_u) = Q_X(L_i)$, and hence $\lambda'_i \equiv \lambda'_{ijk} \simeq \tilde{\mu}_i \equiv \mu_i/\mu$, where the supersymmetry preserving μ parameter is assumed to be of the same order as the supersymmetry breaking soft masses (\tilde{m}). Turning our attention to the $L_i L_j E_k^C$ operator, we notice that when $j = k$, the same argument as above leads to $\lambda_{ijj} \simeq \lambda'_{ijk} \simeq \tilde{\mu}_i$. Thus 39 *a priori* independent L -violating couplings basically boil down to only 6:

$$\tilde{\mu}_i (\simeq \lambda'_{ijk} \simeq \lambda_{ijj}) , \quad \lambda_{123}, \lambda_{132}, \lambda_{231}. \quad (3)$$

Thanks to the flavor symmetry, the L -violating bilinear soft parameters B_i would be aligned to the corresponding superpotential parameters μ_i as well, i.e. $\tilde{B}_i \equiv B_i/\tilde{m}^2 \simeq \tilde{\mu}_i$. It should be noted that when we say that two couplings are related, we mean that they have a common suppression factor ε^{Q_X} . Indeed, there are order-one uncertainties in the actual coefficients of the operators, for which reason we have used a ‘near-equality’ sign in Eq. (3). Now we come to the relative sizes of the L -violating couplings. The suppression would depend on the sum of Q_X charges of the associated lepton fields as a power of ε . More specifically,

$$\tilde{\mu}_i \simeq \tilde{B}_i \simeq \lambda'_i \sim \varepsilon^{Q_X(L_i)}, \quad \lambda_{ijk} \sim \varepsilon^{\{Q_X(L_i)+Q_X(L_j)+Q_X(E_k^C)\}}. \quad (4)$$

Eventually, we shall provide a specific demonstration with $Q = Z_{N_1} \times Z_{N_2}$ [9], which means there are all together 6 charges for the three lepton generations.

Many RpV couplings which are not so strongly constrained may now be related by Eq. (4) to the ones which are severely constrained by experiments. The existing bounds on the individual and product couplings can be found in the reviews [3].

III Neutrino masses and mixing

One of the high points of R -parity violation is that it generates neutrino masses and mixing through a perfectly renormalizable interaction without the need of introducing any right-handed neutrino. This has already been

Our couplings	Related to	Existing limits (Sources)	Refs.
$\tilde{\mu}_i$	$\tilde{\mu}_i, \lambda'_{ijk}, \lambda_{ijj}$	$1.5 \times 10^{-6} (M_{\tilde{\chi}}) [m_{\nu_i}]$	[7]
λ_{123}	λ_{123}	$0.03 (\tilde{\tau}_R) [V_{ud}]$	[4]
λ_{132}	λ_{132}	$0.03 (\tilde{\mu}_R) [R_\tau]$	[4]
λ_{231}	λ_{231}	$0.05 (\tilde{e}_R) [R_\tau]$	[4]

Table 1: The list of the six independent couplings and the standard couplings they are related to by the flavor symmetry X . The three $\tilde{\mu}_i$ couplings are of the same order of magnitude as 36 out of 39 *a priori* independent RpV couplings. A mass of 100 GeV is assumed for the superparticles exchanged in the processes involved. These superparticles are indicated within first bracket right after the bounds (the weak gaugino mass $M_{\tilde{\chi}}$ and the three scalar leptons $\tilde{\ell}_R$). The entries in the square brackets specify the different observables from which origin the bounds originate. Here, $R_\tau = \Gamma(\tau^- \rightarrow \mu^- \bar{\nu}_\mu \nu_\tau) / \Gamma(\mu^- \rightarrow e^- \bar{\nu}_e \nu_\mu)$.

studied at various levels of detail [6, 7, 8]. In this work we will follow the notation of [8]. The neutrino masses, in the basis in which all the sneutrino vacuum expectation values vanish, can be written as

$$\begin{aligned}
m_{ij} \approx & \frac{\cos^2 \beta}{\tilde{m}} \mu_i \mu_j + \frac{g^2}{64\pi^2 \cos^2 \beta} \frac{B_i B_j}{\tilde{m}^3} + \frac{g^2}{64\pi^2 \cos \beta} \frac{\mu_i B_j + \mu_j B_i}{\tilde{m}^2} \\
& + \sum_k \frac{3}{16\pi^2} g m_{d_k} \frac{\mu_i \lambda'_{jkk} + \mu_j \lambda'_{ikk}}{\tilde{m}} + \sum_k \frac{1}{16\pi^2} g m_{e_k} \frac{\mu_i \lambda_{jkk} + \mu_j \lambda_{ikk}}{\tilde{m}} \\
& + \sum_{l,k} \frac{3}{8\pi^2} \lambda'_{ilk} \lambda'_{jkl} \frac{m_{d_l} m_{d_k}}{\tilde{m}_q^2} \mu \tan \beta + \sum_{l,k} \frac{1}{8\pi^2} \lambda_{ilk} \lambda_{jkl} \frac{m_{e_l} m_{e_k}}{\tilde{m}^2} \mu \tan \beta,
\end{aligned} \tag{5}$$

where m_{d_i} (m_{e_i}) denote the masses of the down-type quarks (charged leptons). A comment on the approximations made above is in order. We have denoted the squark masses by \tilde{m}_q and assumed them to be somewhat heavier than a common mass scale \tilde{m} assumed for the sleptons and weak gauginos/Higgsinos. This approximation may be crude but is good enough for our order-of-magnitude estimate of the RpV couplings. In Eq. (5), the first line accounts for the tree level and one loop contributions from bilinear couplings only, the second line represents the one loop contributions from both bilinear and trilinear couplings, while the last line stands for one loop contributions from trilinear couplings only. The possibility of large left-right squark/slepton mixing which may be induced by large $\tan \beta$ ($\equiv v_u/v_d$) has been taken into account in the purely trilinear loop dynamics. The tree level $\mu_i \mu_j$ contribution generates a rank-one mass matrix and therefore yields only one mass eigenvalue. Since, in our case, $B_i, \lambda'_i, \lambda_{ijj}$ are all proportional to μ_i , even after including their contributions the rank-one nature of the mass matrix does not change. What breaks the alignment and yields more non-vanishing eigenvalues is the contribution from the purely trilinear loops involving λ_{ijk} ($i \neq j \neq k$), since these couplings are not aligned with μ_i .

This leaves us with the remaining three couplings, namely, λ_{123} , λ_{132} and λ_{231} , no two indices of which are the same, for generating the second *mandatory* and the third *optional* nonvanishing neutrino masses and the three mixing angles (two large and one small). Note that the existing bounds on λ_{ijk} with $i \neq j \neq k$ are relatively less stringent – see Table 1.

Different low energy processes, especially some lepton flavor violating decays, yield important constraints on trilinear product couplings [12, 13, 14]. Due to the smallness of *most* of the couplings as shown in the first row of Table 1, these constraints are in almost all cases easily satisfied. The bounds emerging from the nonobservation of $K_L^0 \rightarrow \mu \bar{e} / e \bar{\mu}$ [13, 14], namely,

$$\lambda_{ijk} \lambda'_{lmn} < 6.7 \times 10^{-9} m_{\tilde{\nu}_{L3}}^2 / (100 \text{ GeV})^2, \tag{6}$$

with the combinations $(ijk)(lmn) : (312)(312), (312)(321), (321)(312), (321)(321)$, play a crucial rôle in neutrino mass/mixing model building in our scenario, as we shall see later. Due to the specific inter-connections among RpV couplings owing to the flavor symmetry, Eq. (6) leads to the following limits:

$$\lambda_{132} \lambda'_3, \quad \lambda_{231} \lambda'_3 < 6.7 \times 10^{-9} m_{\tilde{\nu}_{L3}}^2 / (100 \text{ GeV})^2. \tag{7}$$

If we set the numerical values of the couplings near their upper limits (see Table 1), they turn out to be large enough to offset the loop suppression factors. The mass matrix entries can then be written with only six RpV couplings as

$$\begin{aligned}
m_{ee} &\approx a\mu_1\mu_1 + \frac{1}{8\pi^2}\lambda_{123}\lambda_{132}\frac{m_\mu m_\tau}{\tilde{m}^2}\mu \tan\beta, \\
m_{e\mu} &\approx a\mu_1\mu_2 + \frac{1}{8\pi^2}\lambda_{123}\lambda_{232}\frac{m_\tau m_\mu}{\tilde{m}^2}\mu \tan\beta + \frac{1}{8\pi^2}\lambda_{213}\lambda_{131}\frac{m_\tau m_e}{\tilde{m}^2}\mu \tan\beta, \\
m_{e\tau} &\approx a\mu_1\mu_3 + \frac{1}{8\pi^2}\lambda_{132}\lambda_{323}\frac{m_\mu m_\tau}{\tilde{m}^2}\mu \tan\beta + \frac{1}{8\pi^2}\lambda_{312}\lambda_{121}\frac{m_\mu m_e}{\tilde{m}^2}\mu \tan\beta, \\
m_{\mu\mu} &\approx a\mu_2\mu_2 + \frac{1}{8\pi^2}\lambda_{231}\lambda_{213}\frac{m_e m_\tau}{\tilde{m}^2}\mu \tan\beta, \\
m_{\mu\tau} &\approx a\mu_2\mu_3 + \frac{1}{8\pi^2}\lambda_{231}\lambda_{313}\frac{m_\tau m_e}{\tilde{m}^2}\mu \tan\beta + \frac{1}{8\pi^2}\lambda_{321}\lambda_{212}\frac{m_\mu m_e}{\tilde{m}^2}\mu \tan\beta, \\
m_{\tau\tau} &\approx a\mu_3\mu_3 + \frac{1}{8\pi^2}\lambda_{312}\lambda_{321}\frac{m_\mu m_e}{\tilde{m}^2}\mu \tan\beta,
\end{aligned} \tag{8}$$

with

$$a = \frac{\cos^2\beta}{\tilde{m}} + \sum_k \frac{3gm_{d_k}}{8\pi^2\tilde{m}^2} + \sum_k \frac{gm_{e_k}}{8\pi^2\tilde{m}^2} + \sum_{k,l} \frac{3m_{d_l}m_{d_k}}{8\pi^2\mu\tilde{m}_q^2} \tan\beta. \tag{9}$$

With this mass matrix we try to reproduce the neutrino oscillation data, namely, the two mass-squared differences (Δm_{21}^2 and Δm_{31}^2) and the three mixing angles (θ_{12} , θ_{23} and θ_{13}). For simplicity we assume that all the phases in the neutrino mixing matrix are zero. Since neutrino oscillation analysis probes only the mass-squared differences and not their absolute values, we need to assume the hierarchy (normal/inverted) of the masses and the size of the smallest eigenvalue to fix the other two masses. There is no lower limit on the smallest neutrino mass eigenvalue, it can still be zero.

We take the best fit values of the neutrino mass-squared differences from [15]: $\Delta m_{21}^2 = 7.59 \times 10^{-5} \text{ eV}^2$, $\Delta m_{31}^2(\text{IH}) = -2.40 \times 10^{-3} \text{ eV}^2$, $\Delta m_{31}^2(\text{NH}) = 2.51 \times 10^{-3} \text{ eV}^2$. Above, NH stands for ‘normal hierarchy’ and IH stands for ‘inverted hierarchy’ of neutrino masses. The two mixing angles θ_{12} and θ_{23} are set to their TBM values (using best fit values instead does not lead to significant changes). Very recently, two long-baseline accelerator experiments T2K and MINOS, both probing $\nu_\mu \rightarrow \nu_e$ appearance, have reported, for the first time, a non-zero measurement of θ_{13} . T2K has observed 6 electron-like events against an estimated background of 1.5, thus discarding $\theta_{13} = 0$ at the level of 2.5σ [16]. The MINOS experiment observes 62 electron-like events against an expected 49, thus disfavoring $\theta_{13} = 0$ at 1.5σ [17]. A new global fit suggests $\sin^2\theta_{13} = 0.021(0.025) \pm 0.007$ with old (new) reactor fluxes [18]. The central value corresponds to $\theta_{13} \approx 9^\circ$. A second global fit can be found in [19]. Both indicate $\theta_{13} > 0$ with a significance of about 3σ . We shall see below that in our scenario whether θ_{13} is vanishing or non-vanishing plays an important rôle in predicting whether neutrino mass hierarchy is *normal* ($\Delta m^2 \equiv m_3^2 - 0.5(m_2^2 + m_1^2) > 0$) or *inverted* ($\Delta m^2 < 0$). Since none of the two experiments has so far conclusively established a nonzero value of θ_{13} , we take both the paradigms, namely, $\theta_{13} = 0$ and $\theta_{13} \neq 0$, and study what do they imply on the choices of RpV parameters and whether we can predict the nature of mass hierarchy.

IV Tri-bimaximal mixing

The TBM structure immediately implies that $|m_{e\mu}| = |m_{e\tau}|$ and $|m_{\mu\mu}| = |m_{\tau\tau}|$, regardless of whether the lightest mass eigenvalue is vanishing or not, and also irrespective of whether the neutrino mass hierarchy is normal or inverted. For our couplings this can be comfortably realized by setting $|\lambda_{123}| = |\lambda_{132}|$ and $|\mu_2| = |\mu_3|$ – see Eq. (8). This means that we can parametrize the mass matrix with four independent RpV parameters instead of six, which of course improves the predictivity of the model. Dropping the terms in the loop contribution

proportional to the electron mass, we obtain

$$m_{e\mu} \approx m_{e\tau} \approx a\mu_1\mu_2 - \frac{1}{8\pi^2}\lambda_{123}\mu_3 \frac{m_\tau m_\mu}{\tilde{m}^2} \tan\beta. \quad (10)$$

Clearly, under this situation, the absolute values for the tree-level contributions to $m_{\mu\mu} \sim a\mu_2\mu_2$, $m_{\mu\tau} \sim a\mu_2\mu_3$ and $m_{\tau\tau} \sim a\mu_3\mu_3$ are the same. Setting all CP-violating phases to zero, the TBM mixing matrix takes the form [20]

$$U_{\text{TBM}} = \begin{pmatrix} \sqrt{\frac{2}{3}} & \frac{1}{\sqrt{3}} & 0 \\ -\frac{1}{\sqrt{6}} & \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{6}} & -\frac{1}{\sqrt{3}} & \frac{1}{\sqrt{2}} \end{pmatrix}. \quad (11)$$

To fix the numerical values of the mass matrix from $m = U_{\text{TBM}}^T \times \text{diag}(m_1, m_2, m_3) \times U_{\text{TBM}}$, all we need to decide is the mass hierarchy (normal or inverted) and the smallest mass eigenvalue.

Inverted hierarchy: We first consider the case of inverted hierarchy with $m_3 = 0$. This choice additionally demands $m_{\mu\tau} = -m_{\mu\mu}$. One obtains

$$m = \begin{pmatrix} 4.92 \times 10^{-2} & 2.56 \times 10^{-4} & -2.56 \times 10^{-4} \\ 2.56 \times 10^{-4} & 2.47 \times 10^{-2} & -2.47 \times 10^{-2} \\ -2.56 \times 10^{-4} & -2.47 \times 10^{-2} & 2.47 \times 10^{-2} \end{pmatrix} \text{ eV (IH, TBM, } m_3 = 0). \quad (12)$$

By setting $\mu_2 = -\mu_3$ and keeping $\lambda_{231} \lesssim \lambda_{123}$, we obtain a rough analytical solution using Eqs. (4) and (8):

$$|\mu_2| = |\mu_3| \approx \sqrt{a^{-1}m_{\mu\mu}}, \quad \lambda_{123} \approx \sqrt{\frac{4\pi^2 m_{ee} \tilde{m}^2}{m_\mu m_\tau \mu \tan\beta}}, \quad \mu_1 \approx \frac{m_{e\mu} + \mu_2 \lambda_{123} m_\tau m_\mu \tan\beta / (8\pi^2 \tilde{m}^2)}{(a\mu_2)}. \quad (13)$$

Putting $\tilde{m} = \mu = 100 \text{ GeV}$ and $\tilde{m}_q = 300 \text{ GeV}$ in Eq. (13) we obtain a solution (with $\mu_2 = -\mu_3$, and $\lambda_{132} = -\lambda_{123}$)

$$\tilde{\mu}_1 = 1.9 \times 10^{-8}, \quad \tilde{\mu}_2 = -4.7 \times 10^{-6}, \quad \lambda_{231} \sim 10^{-4}, \quad \lambda_{123} = -3.2 \times 10^{-4} \text{ for } \tan\beta = 10. \quad (14)$$

To illustrate how this coupling pattern can arise from a flavor symmetry we are providing an exemplary flavor group for this case. However, since this choice is not necessarily unique and our conclusions do not depend on the specific flavor group, we omit this exercise for the other scenarios. In this case, the required relative suppression can be reproduced by a $Q_X = Z_4 \times Z_8$ family symmetry with a breaking parameter ε . The necessary charge assignments are given by,

$$Q_B(L_1) = (2, 5), \quad Q_B(L_2) = (0, 5), \quad Q_B(L_3) = (3, 2), \quad (15)$$

which imply

$$Q_B(L_1 L_2 E_3^C) = (3, 0), \quad Q_B(L_1 L_3 E_2^C) = (1, 2), \quad Q_B(L_2 L_3 E_1^C) = (1, 2). \quad (16)$$

These assignments lead exactly to the required suppression of the couplings with $\varepsilon \approx 0.1$ as

$$\mu_2(= \mu_3) \sim \varepsilon^5, \quad \mu_1 \sim \varepsilon^7, \quad \lambda_{123}(= \lambda_{132}) \sim \varepsilon^3, \quad \lambda_{231} \sim \varepsilon^3. \quad (17)$$

In the above example, the near equality of the magnitude of the entries in the $\mu - \tau$ block is ensured by saturating them with the tree level contributions, while keeping the loop contributions suppressed. If, within the TBM framework, we now consider m_3 to be slightly above zero, then $m_{\mu\mu} = m_{\tau\tau} > |m_{\mu\tau}|$. To obtain $m_3 = 0.001 \text{ eV}$ with $\tilde{m} = \mu = 100 \text{ GeV}$, we need $\tilde{\mu}_1 = 1.9 \times 10^{-8}$, $\tilde{\mu}_2 = -4.6 \times 10^{-6}$, $\tilde{\mu}_3 = 4.7 \times 10^{-6}$, $\lambda_{123} = -3.1 \times 10^{-4}$, $\lambda_{132} = -3.3 \times 10^{-4}$, $\lambda_{231} = 2.7 \times 10^{-3}$. We should note two important things: (i) These choices imply $\lambda_{231}\lambda_3' = 1.3 \times 10^{-8}$, which mildly overshoots the $K_L \rightarrow \mu e$ bound as shown in Eq. (7). If we increase m_3 further, the disagreement with the K_L bounds deepens. (ii) The ‘four parameter’ scenario with

$|\mu_2| = |\mu_3|$ and $|\lambda_{123} = \lambda_{132}|$ is not compatible with a non-vanishing absolute neutrino mass scale, i.e. we cannot fit the data assuming these ‘equalities’ with $m_3 > 0$, because of the hierarchical nature of the charged lepton masses which appear in Eq. (8).

Normal hierarchy: We now consider normal hierarchy of neutrino masses. In this case the smallest mass eigenvalue is m_1 . Within the TBM structure if we keep $m_1 = 0$, it follows that $m_{\mu\mu} = m_{\tau\tau} > |m_{\mu\tau}|$. The numerical values of the mass matrix entries are

$$m = \begin{pmatrix} 2.90 \times 10^{-3} & 2.90 \times 10^{-3} & -2.90 \times 10^{-3} \\ 2.90 \times 10^{-3} & 2.80 \times 10^{-2} & 2.21 \times 10^{-2} \\ -2.90 \times 10^{-3} & 2.21 \times 10^{-2} & 2.80 \times 10^{-2} \end{pmatrix} \text{ eV (NH, TBM, } m_1 = 0). \quad (18)$$

The couplings needed to fit these entries are $\tilde{\mu}_1 = -5.2 \times 10^{-7}$, $\tilde{\mu}_2 = 3.9 \times 10^{-6}$, $\tilde{\mu}_3 = 5.0 \times 10^{-6}$, $\lambda_{123} = -4.4 \times 10^{-3}$, $\lambda_{132} = -1.2 \times 10^{-6}$, $\lambda_{231} = 1.0 \times 10^{-3}$. Although we are within the $K_L \rightarrow \mu e$ bound, the requirement $m_{e\mu} = -m_{e\tau}$ is realized quite differently. The relative signs of the tree-level couplings invariably imply $m_{e\mu}^{\text{tree}} \approx +m_{e\tau}^{\text{tree}}$. This difference between the experimental requirement and the tree-level contribution cannot be resolved, even keeping in mind that signs of the RpV couplings can be chosen at will and also each neutrino field can be redefined to absorb a sign. Therefore, a sign adjustment for one of the entries ($e\mu$) via a large loop contribution is needed, while the loop contribution to the other one ($e\tau$) becomes negligible. This is reflected in the large hierarchy between λ_{123} and λ_{132} . We recall that such a sign adjustment was not required in the case of inverted hierarchy (TBM, $m_3 = 0$). If we now increase the value of m_1 (from zero) and try to fit normal hierarchy within the TBM framework, the $K_L \rightarrow \mu e$ bound haunts us like in the case of inverted hierarchy with $m_3 > 0$. Therefore, our most robust prediction is the tight constraint for the smallest mass eigenvalue. Thus, inverted hierarchy can be fit with four parameters, while normal hierarchy requires six parameters and a sign altering large loop correction.

V Non-zero θ_{13}

In view of the recent T2K data which measures non-vanishing θ_{13} , we study how flexible we are to accommodate normal or inverted hierarchies while keeping θ_{13} close to its central value of 9.0° . Unlike in the case of TBM which guarantees $|m_{e\mu}| = |m_{e\tau}|$ and $m_{\mu\mu} = m_{\tau\tau}$, it is not possible to fit the data with 4 parameters when $\theta_{13} \neq 0$.

Inverted hierarchy: First we consider the case $m_3 = 0$. The numerical entries of the mass matrix are given by

$$m = \begin{pmatrix} 4.80 \times 10^{-2} & -5.13 \times 10^{-3} & -5.63 \times 10^{-3} \\ -5.13 \times 10^{-3} & 2.53 \times 10^{-2} & -2.41 \times 10^{-2} \\ -5.63 \times 10^{-3} & -2.41 \times 10^{-2} & 2.54 \times 10^{-2} \end{pmatrix} \text{ eV (IH, } \theta_{13} = 9.0^\circ, m_3 = 0). \quad (19)$$

This can be fit with $\tilde{\mu}_1 = -1.1 \times 10^{-6}$, $\tilde{\mu}_2 = -4.5 \times 10^{-6}$, $\tilde{\mu}_3 = 4.8 \times 10^{-6}$, $\lambda_{123} = 9.3 \times 10^{-3}$, $\lambda_{132} = 1.1 \times 10^{-5}$, $\lambda_{231} = -1.1 \times 10^{-4}$. Two things are worth noting: (i) The magnitudes of λ_{123} and λ_{132} are separated by nearly three orders, while they assumed identical numerical values in the case of TBM. (ii) The tree-level contribution to $m_{e\mu}$ has the wrong sign like in the case of NH with $\theta_{13}=0$. Again a large sign adjusting loop contribution is needed to be in agreement with the experimental data. If we now increase the value of m_3 , the required magnitude for λ_{231} becomes larger, and eventually beyond $m_3 = 0.01$ eV the $K_L \rightarrow \mu e$ bound overshoots.

Normal hierarchy: For $m_1 = 0$, the mass matrix entries are given by

$$m = \begin{pmatrix} 4.06 \times 10^{-3} & 8.02 \times 10^{-3} & 2.29 \times 10^{-3} \\ 8.02 \times 10^{-3} & 2.67 \times 10^{-2} & 2.16 \times 10^{-2} \\ 2.29 \times 10^{-3} & 2.16 \times 10^{-2} & 2.80 \times 10^{-2} \end{pmatrix} \text{ eV (NH, } \theta_{13} = 9.0^\circ, m_1 = 0). \quad (20)$$

This can be reproduced with $\tilde{\mu}_1 = 4.1 \times 10^{-7}$, $\tilde{\mu}_2 = 3.8 \times 10^{-6}$, $\tilde{\mu}_3 = 5.0 \times 10^{-6}$, $\lambda_{123} = -5.3 \times 10^{-3}$, $\lambda_{132} = -1.5 \times 10^{-6}$, $\lambda_{231} = 8.3 \times 10^{-4}$. Note that λ_{231} is small enough to satisfy the $K_L \rightarrow \mu e$ bound.

Contrary to the case of inverted hierarchy, now no large sign-flipping correction for $m_{e\mu}$ is needed. However, the difference between the values of $m_{e\mu}$ and $m_{e\tau}$ still leads to a hierarchy in the λ -couplings. Just like in the previous cases, the $K_L \rightarrow \mu e$ bound begins to be relevant as soon as m_1 increases to around 0.005 eV (which requires $\lambda_{231} = 1.5 \times 10^{-3}$). The main conclusion for non-zero θ_{13} is again that the smallest mass eigenvalue is required to be almost vanishing in both hierarchies. But contrary to the TBM case, now IH requires a sign adjustment, while NH does not.

VI Collider signatures

The LHC signatures of the λ_{ijk} couplings have recently been explored in [21]. In our scenario, only three couplings λ_{ijk} ($i \neq j \neq k$) are relatively large ($10^{-3} - 10^{-4}$), the rest are of order 10^{-6} . The large couplings are small enough to make sure that the RpV vertex is numerically relevant only at the end of a supersymmetry cascade when the lightest neutralino decays via a λ_{ijk} interaction, $\tilde{\chi}_1^0 \rightarrow l^\pm l^\mp \nu$. The λ_{ijk} couplings thus give rise to $l_i l_k$ or $l_j l_k$ final states plus missing energy. Depending on the numerical values of the corresponding λ_{ijk} couplings the branching ratios into the $l_i l_k$ or $l_j l_k$ channel will scale as $|\lambda_{ijk}|^2$. Thus both invariant mass distributions and number counting of the final state leptons should be a part of the search method. However, other decay channels like $\tilde{\chi}_1^0 \rightarrow W^\pm l^\mp$ and $\tilde{\chi}_1^0 \rightarrow Z \nu$ are available due to the presence of the bilinear couplings. Their role has been investigated in detail in [22]. Therefore, a detailed study of neutralino decays is important to test this and other RpV models and differentiate between them. The non-observation of an excess in four lepton events at CMS and ATLAS so far indicates a somewhat heavier squark mass scale than the one we choose. However, scaling the slepton masses accordingly, this will not lead to any significant changes related to our work.

VII Conclusions

In this paper we have studied a generic and simple flavor model which reduces the number of independent couplings from 39 to 6, i.e. μ_i ($i = 1, 2, 3$), λ_{123} , λ_{132} and λ_{231} . This results in an extremely predictive framework, which can reproduce the correct neutrino masses and mixings while satisfying all other low energy bounds.

In its simplest realization the scenario leads to a four parameter model with exact tri-bimaximal mixing and prefers inverse hierarchy, for which a specific flavor model, viz $Z_4 \times Z_8$, has been proposed. A non-vanishing mixing angle θ_{13} can be accommodated in a six parameter realization.

A general prediction of all possible realizations is an almost vanishing absolute mass scale for neutrinos, i.e. an essentially massless lightest neutrino. This feature is tightly related to the non-observation of $K_L \rightarrow \mu e$ which affects many important coupling products in this framework. As a consequence, any positive signal in one of the upcoming neutrinoless double beta decay experiments would imply an inverted neutrino mass hierarchy, since for the combination of normal hierarchy and an almost vanishing absolute mass scale, the resulting $|m_{ee}|$ is beyond their sensitivity. In other words, if a conclusive evidence of nonzero θ_{13} is established, then our scenario would be able to accommodate a positive signal of neutrinoless double beta decay only at the expense of large sign-flipping correction to one of the off-diagonal elements of the mass matrix. Moreover, the flavor structure proposed here can lead to specific decays of a neutralino LSP at the LHC.

Acknowledgments:

We thank W.Porod for valuable suggestions. This work was supported by DAAD-DST PPP Grant No. D/08/04933 and DST-DAAD project No. INT/DAAD/P-181/2008. G.B. acknowledges the *Gambrinus*

grant and hospitality at T.U. Dortmund during a part of the collaboration. D.P. and H.P. were supported by DFG Grant No. PA 803/5-1, the Physics at the Terascale Helmholtz Alliance Working Group: Neutrino masses and Lepton Flavor Violation at the LHC, and they acknowledge hospitality at the Saha Institute of Nuclear Physics, Kolkata, during a part of this collaboration.

References

- [1] G. R. Farrar and P. Fayet, *Phys. Lett. B* **76** (1978) 575; S. Weinberg, *Phys. Rev. D* **26** (1982) 287; N. Sakai and T. Yanagida, *Nucl. Phys. B* **197** (1982) 533; C. S. Aulakh and R. N. Mohapatra, *Phys. Lett. B* **119** (1982) 136.
- [2] A. Santamaria and J. W. F. Valle, *Phys. Rev. Lett.* **60** (1988) 397; A. Santamaria and J. W. F. Valle, *Phys. Lett. B* **195** (1987) 423; A. Santamaria and J. W. F. Valle, *Phys. Rev. D* **39** (1989) 1780; J. C. Romao and J. W. F. Valle, *Nucl. Phys. B* **381** (1992) 87; J. C. Romao, C. A. Santos and J. W. F. Valle, *Phys. Lett. B* **288** (1992) 311.
- [3] For reviews see, G. Bhattacharyya, *Nucl. Phys. Proc. Suppl.* **52A** (1997) 83 [arXiv:hep-ph/9608415]; G. Bhattacharyya, arXiv:hep-ph/9709395; H. K. Dreiner, arXiv:hep-ph/9707435; M. Chemtob, *Prog. Part. Nucl. Phys.* **54** (2005) 71 [arXiv:hep-ph/0406029]; R. Barbier *et al.*, *Phys. Rept.* **420** (2005) 1 [arXiv:hep-ph/0406039].
- [4] For a very recent update on single RpV couplings, see Y. Kao and T. Takeuchi, arXiv:0910.4980 [hep-ph].
- [5] Sakai and Yanagida in [1]; A. Y. Smirnov and F. Vissani, *Phys. Lett. B* **380** (1996) 317 [arXiv:hep-ph/9601387]; G. Bhattacharyya and P. B. Pal, *Phys. Rev. D* **59** (1999) 097701 [arXiv:hep-ph/9809493]; G. Bhattacharyya and P. B. Pal, *Phys. Lett. B* **439** (1998) 81 [arXiv:hep-ph/9806214].
- [6] L. J. Hall and M. Suzuki, *Nucl. Phys. B* **231** (1984) 419; M. Nowakowski and A. Pilaftsis, *Nucl. Phys. B* **461** (1996) 19 [arXiv:hep-ph/9508271]; S. Rakshit, G. Bhattacharyya and A. Raychaudhuri, *Phys. Rev. D* **59** (1999) 091701 [arXiv:hep-ph/9811500]; G. Bhattacharyya, H. V. Klapdor-Kleingrothaus and H. Päs, *Phys. Lett. B* **463**, 77 (1999) [arXiv:hep-ph/9907432]; D. E. Kaplan and A. E. Nelson, *JHEP* **0001** (2000) 033 [arXiv:hep-ph/9901254]; M. Hirsch, M. A. Diaz, W. Porod, J. C. Romao and J. W. F. Valle, *Phys. Rev. D* **62** (2000) 113008 [Erratum-ibid. *D* **65** (2002) 119901] [arXiv:hep-ph/0004115]; M. A. Diaz, M. Hirsch, W. Porod, J. C. Romao and J. W. F. Valle, *Phys. Rev. D* **68** (2003) 013009 [Erratum-ibid. *D* **71** (2005) 059904] [arXiv:hep-ph/0302021]; S. Davidson and M. Losada, *Phys. Rev. D* **65** (2002) 075025 [arXiv:hep-ph/0010325]; A. Dedes, S. Rimmer and J. Rosiek, *JHEP* **0608** (2006) 005 [arXiv:hep-ph/0603225]; A. S. Joshipura and M. Nowakowski, *Phys. Rev. D* **51** (1995) 2421 [arXiv:hep-ph/9408224]; R. Hempfling, *Nucl. Phys. B* **478** (1996) 3 [arXiv:hep-ph/9511288]. M. Hirsch, H. V. Klapdor-Kleingrothaus and S. G. Kovalenko, *Phys. Lett. B* **398** (1997) 311 [arXiv:hep-ph/9701253]. Y. Grossman and H. E. Haber, *Phys. Rev. Lett.* **78** (1997) 3438 [arXiv:hep-ph/9702421]. Y. Grossman and H. E. Haber, *Phys. Rev. D* **59** (1999) 093008 [arXiv:hep-ph/9810536]. Neutrino mass calculation induced by R -parity violation is now incorporated in SOFTSUSY code, see B. C. Allanach, C. H. Kom, M. Hanussek, [arXiv:1109.3735 [hep-ph]].
- [7] A. Abada, S. Davidson and M. Losada, *Phys. Rev. D* **65** (2002) 075010 [arXiv:hep-ph/0111332]; S. Davidson and M. Losada, *JHEP* **0005** (2000) 021 [arXiv:hep-ph/0005080]; A. Abada, G. Bhattacharyya and M. Losada, *Phys. Rev. D* **66** (2002) 071701 [arXiv:hep-ph/0208009].
- [8] Y. Grossman and S. Rakshit, *Phys. Rev. D* **69** (2004) 093002 [arXiv:hep-ph/0311310].
- [9] T. Banks, Y. Grossman, E. Nardi and Y. Nir, *Phys. Rev. D* **52** (1995) 5319 [arXiv:hep-ph/9505248].
- [10] G. Bhattacharyya, *Phys. Rev. D* **57** (1998) 3944 [arXiv:hep-ph/9707297]; Y. Kajiyama, *JHEP* **0704** (2007) 007 [arXiv:hep-ph/0702056]; J. R. Ellis, S. Lola and G. G. Ross, *Nucl. Phys. B* **526** (1998) 115 [arXiv:hep-ph/9803308].

- [11] H. K. Dreiner, J. Soo Kim, M. Thormeier, [arXiv:0711.4315 [hep-ph]].
- [12] G. Bhattacharyya, K. B. Chatterjee and S. Nandi, Nucl. Phys. B **831** (2010) 344 [arXiv:0911.3811 [hep-ph]].
- [13] H. K. Dreiner, M. Kramer and B. O’Leary, Phys. Rev. D **75**, 114016 (2007) [arXiv:hep-ph/0612278].
- [14] H. K. Dreiner, G. Polesello and M. Thormeier, Phys. Rev. D **65** (2002) 115006 [arXiv:hep-ph/0112228].
- [15] M. C. Gonzalez-Garcia, M. Maltoni and J. Salvado, JHEP **1004** (2010) 056 [arXiv:1001.4524 [hep-ph]].
- [16] K. Abe *et al.* [T2K Collaboration], arXiv:1106.2822 [hep-ex].
- [17] L. Whitehead [MINOS Collaboration], “Recent results from MINOS”, Joint Experimental-Theoretical Seminar (24 June 2011, fermilab, USA).
- [18] G. L. Fogli, E. Lisi, A. Marrone, A. Palazzo and A. M. Rotunno, arXiv:1106.6028 [hep-ph].
- [19] T. Schwetz, M. Tortola, J. W. F. Valle, [arXiv:1108.1376 [hep-ph]].
- [20] P. F. Harrison, D. H. Perkins, W. G. Scott, Phys. Lett. **B530** (2002) 167. [hep-ph/0202074].
- [21] N. -E. Bomark, D. Choudhury, S. Lola, P. Osland, JHEP **1107** (2011) 070. [arXiv:1105.4022 [hep-ph]].
- [22] W. Porod, M. Hirsch, J. Romao, J. W. F. Valle, Phys. Rev. **D63** (2001) 115004. [hep-ph/0011248].